

NSAIDS

- Choline and magnesium salicylates (CMT, Tricosal, Trilisate)
- Choline salicylate (Arthropan)
- Diclofenac potassium (Cataflam)
- Diclofenac sodium (Voltaren, Voltaren XR)
- Diclofenac sodium with misoprostol (Arthrotec)
- Diflunisal (Dolobid)
- Etodolac (Lodine, Lodine XL)
- Fenoprofen calcium (Nalfon)
- Flurbiprofen (Ansaid)
- **Ibuprofen (Advil, Motrin, Motrin IB, Nuprin)**
- Indomethacin (Indocin, Indocin SR)
- Ketoprofen (Aaron, Orudis, Orudis KT, Oruvail)
- Magnesium salicylate (Arthritab, Bayer Select, Doan's Pills, Magan, Mobidin, Mobogesic)
- Meclofenamate sodium (Meclomen)
- Mefenamic acid (Ponstel)
- Meloxicam (Mobic)
- Nabumetone (Relafen)
- Naproxen (Naprosyn, Naprelan)
- Naproxen sodium (Aleve, Anaprox)
- Oxaprozin (Daypro)
- Piroxicam (Feldene)
- Rofecoxib (Vioxx)
- Salsalate (Amigesic, Anaflex 750, Disalcid, Marthritic, Mono-Gesic, Salflex, Salsitab)
- Sodium salicylate (various generics)
- Sulindac (Clinoril)
- Tolmetin sodium (Tolectin)
- Valdecoxib (Bextra)

Note: Some products, such as Excedrin, are combination drugs (Excedrin is acetaminophen, aspirin, and caffeine).

BLOOD THINNERS

- Aggrenox (ASA + Dipyridamole)
- Arixtra (fondaparinux sodium)
- **Aspirin (Anacin, Ascriptin, Bayer, Bufferin, Ecotrin, Excedrin)**
- Coumadin (Warfarin)
- Fragmin (Dalteparin sodium)
- Heparin (Heparin sodium)
- Lovenox (Enoxaparin sodium)
- Persantine (Dipyridamole)
- Plavix (Clopidogrel)
- Ticlid (Ticlopidine HCL)

PLEASE NOTE THAT BLOOD THINNERS MAY BE RESTARTED TWO (2) WEEKS AFTER SURGERY UNLESS OTHERWISE INDICATED BY THE PHYSICIAN THAT PRESCRIBED THEM.